

SPECIFIC USE INSTRUCTIONS

- Apply Pinpoint™ to dry turfgrass or after mowing
- Allow turf to dry before irrigation when targeting foliar disease
- For control of soil-borne disease, Pinpoint may be watered in immediately
- For foliar disease, use 8L spray solution per 100 m²
- For Dollar spot control, repeat at a 14-21 day interval
- · Application should not be made if rain is expected within two hours
- One 1.77 L jug treats 2.01 1.79 Ha (4.9 4.4 ac)

CHEMICAL CLASS	Group 11
APPLICATION RATE	Chart below
RE-ENTRY INTERVAL	Once product has dried
PACKAGING	1.77 L jug (4 per case)

DISEASES CONTROLLED

- Dollar spot
- Brown patch
- Fairy ring
- Rust (suppression)
- Take-all patch

RECOMMENDED GRASS SPECIES

- Annual bluegrass
- Bentgrass
- Fescue
- Kentucky bluegrass
- Perennial ryegrass

BENEFITS

- Outstanding Dollar spot control
- Preventive and early curative activity
- · Excellent treated turf health and quality
- Tank-mixable with most other products
- Great tank-mix partner with Tourney®
- Pinpoint Fungicide was developed for use without an adjuvant

REGISTERED PESTS	USE RATES
Dollar spot (Sclerotinia homoeocarpa)	NEW 880 - 986 ml/ha
Brown patch (Rhizoctonia solani)	986 ml/ha
Fairy ring	986 ml/ha
Rust (<i>Puccinia grammis</i>) - suppression	986 ml/ha
Take-all patch (Gaeumannomyces graminis)	986 ml/ha

Use rate is dependent on target pest as well as pest stage and severity



