

Effective, economical weed control for turf,  
including golf courses and sod farms.



# Trillion® OS

## CHALLENGE

Sod farms, golf courses and other turf growers struggle to find the right solution to tackle tough weed problems. Too many brands and products fall short of providing a winning combination of effective weed control at a great price.

## NUFARM SOLUTION

Trillion® OS turf herbicide delivers the hard-hitting answer to tough turf weeds including dandelion, chickweed, plantain and creeping Charlie. The familiar Group 4 formulation provides the performance and price you need in easy-to-handle jugs or special order totes.

## REGISTERED USES

Turfgrass including golf courses, sod farms, and municipal, industrial and residential turf

WEEDS CONTROLLED	APPLICATION RATES & NOTES
Bedstraw	<b>TURF (OTHER THAN BENTGRASS):</b> On established turf, apply at the rate of 60 mL per 100 m <sup>2</sup> in 10 L of water.
Black medic (yellow clover)	For larger areas apply at the rate of 5.5 L per hectare in 300 L of water.
Buttercup	
Chickweed	
Clover	
Daisy species (English and ground)	<b>FOR BENTGRASS:</b> Apply at a half rate – 30 mL per 100 m <sup>2</sup> in May or late August and September. Use 20 L of water per 100 m <sup>2</sup> . Slight yellowing will occur but will disappear after about 1 week. Do not overdose bentgrass as damage or kill may occur. Make sure correct rate is applied to the whole area treated. When spraying bentgrass use a 1 nozzle sprayer. Make sure spray application does not overlap or excess material will be applied to overlapped areas and damage is more likely to occur.
Dandelion	
Devil's paint brush	
Ground ivy (creeping Charlie)	
Heal-all	
Ironweed	
Knotweed	
Plantain	
Poison ivy	
Ragweed	
Shepherd's-purse	

## PERFORMANCE TIPS

- For best results apply when weeds are growing quickly
- If weed populations do not warrant a broadcast application, consider spot treatments that target only weedy areas
- Grass seed can be safely sown 1-2 weeks after application at recommended rate
- Do not water turf within 24 hours of application
- Do not apply to newly seeded lawns until after third mowing
- Do not freeze

## LEGEND



## ACTIVE INGREDIENT

- 2,4-D (190 g a.e./L),  
mecoprop-p (100 g a.e./L),  
dicamba (18 g a.e./L)

## FORMULATION

- liquid

## CHEMICAL FAMILY

- phenoxy-carboxylic acid

## PACKAGE SIZE

- 2 x 10 L jug, 155 L drum, 1000 L tote
- 1 jug treats 4.5 acres (at 5.5 L/ha)

## RAINFALL

- do not spray if rain is forecast

## APPLICATION / SEASON

- up to 2 broadcast applications

## SURFACTANT NEEDED

- no

<b>PCP #</b>	<b>WSSA GROUP #</b>
• 35733	• 4

